

Top-down archaeology: High resolution satellite images of Rapa Nui on Google Earth™*

Terry L. Hunt

Department of Anthropology, University of Hawai'i

Carl P. Lipo

Department of Anthropology and IIRMES, California State University Long Beach

A CONSERVATION ETHIC

The archaeological record is, by its very nature, nonrenewable. Each aspect of the record comprises unique information, the sum of which provides the only testament for vast amounts of human history in space and time. The recognition of the nonrenewable character of the archaeological record has led to the compelling rationale for a "conservation ethic" (Lipo 1974).

A conservation ethic demands that we minimize damage to the archaeological record whenever possible for two different, but complementary reasons (Dunnell and Dancey 1978; Dunnell 1984; Moratto and Kelly 1977). First, we must endeavor to preserve cultural heritage for its own sake, since it has intrinsic and diverse values to living populations. Such heritage has cultural, symbolic, political, and aesthetic values, among others. Second, preservation of archaeological phenomena is a scientific imperative, since such phenomena comprise the only empirical record of our past that is subject to scientific explanation. Thus as archaeologists we must seek to preserve the record even in the course of our continuing research. As students of archaeology learn from the outset, archaeological research is often destructive – we sometimes destroy what we study and seek to preserve – like the metaphor of tearing the pages from a book as we read it. Despite this paradox, archaeologists must be fervent advocates for preservation.

In continuing research of the archaeological record the conservation ethic should guide our choices of data generation techniques. While the "standard" techniques of excavation and survey are just two sets of strategies that potentially provide us information, we need to document the archaeological record in ways that best address our research questions. Combining research interests and a conservation ethic, our field strategies comprise the *cheapest* (i.e., most efficient) means for generating observations of archaeological phenomena, which in turn employs means that are usually the *least damaging* to the record. Choosing field research techniques should include as many kinds of observations as possible since the record has potentially many artificial characteristics expressed in visible light, electromagnetic spectra, as well as variations in chemical and physical form resulting from human behavior and expressed at scales ranging from the ion to the landscape. We should acquire

and overlay as many classes of information as possible, with minimal impacts to the archaeological record, employing iterative sampling designs generating data until reaching redundancy in specified classes of information at levels of precision required to resolve problems of interest. For scientific archaeology, Dunnell (1982:7) has described the methodological dimensions of empirical sufficiency (i.e., Are units of meaning measurable in the empirical realm?) and their relation to tolerance limits (How close is close enough in our measurements?); see also Hunt et al. 2001. In concert with dynamic sufficiency (i.e., completeness of theory), our efforts should focus on decreasing the gaps between explanatory accounts (i.e., models and hypotheses) and our particular case (Dunnell 1982; Hunt et al. 2001).

"TOP-DOWN" ARCHAEOLOGY

The organization of the dizzying array of possible archaeological observations into a research management strategy is accomplished by taking a "top-down" approach (Figure 1). In this way, we begin our work with inexpensive and high-

<i>Technique</i>	<i>Scale</i>	<i>Invasiveness</i>
Satellite Image	Large	Minimum
Aerial Photography	↓	↓
Kite Photography		
Pedestrian Survey		
Geophysical Survey		
Surface Mapping		
Auger holes	Small	Maximum
Excavation		

Figure 1. A top-down field research model for archaeology emphasizing the conservation ethic.

* This paper has been peer-reviewed. Paper received 12 August 2007; Accepted 13 January 2008; Revised 28 January 2008.

resolution satellite imagery to determine the distribution of alterations from human actions across a landscape. We then could use publicly-available infrared, sub-meter resolution low-level aerial photographs available for many years and different seasons to map topography, soil and vegetation differences or to examine areas in their historic context. Based on hypotheses about the relationship of the composition of the archaeological record with the landscape generated from the satellite and aerial images, we can then zoom in on areas with low-level aerial photographs available from kite- or balloon-based platforms to document structures and objects at the scale of portable artifacts. For example, we could ask how patterns visible in small-scale imagery might be areas that are compositionally distinct as a consequence of cultural debris deposited on the surface.

Today, using guidance from satellite and aerial images, we might expand our sensory picture of the target area through intensive ground-based surveys and surface mapping to generate data for the spatial distribution of sub-meter scale artifacts and the details of features. This work might be followed with geophysical studies of images in predetermined areas in order to map compositional information about subsurface and near-surface deposits. Targeted areas could also be characterized using geophysical methods such as resistivity and ground penetrating radar or any of the other near-surface remote sensing techniques that are currently available. Each of these techniques provides a means of resolving *structure* (i.e., patterns of variability) in the archaeological record. Subsequent work might layer information obtained by inexpensive small-hole excavations (i.e., coring) that can be used to acquire micro-artifact samples as well as smaller sediment and ion-sized artifact information (e.g., “artificial” chemical concentrations such as phosphorous). These classes of data at combined scales form a robust understanding of the archaeological record in the context of the landscape upon which it is distributed.

RAPA NUI

On Rapa Nui, some of the scales of inquiry have seen more use than others. Landscape approaches have tended to dominate research practices (e.g., McCoy 1976; Cristino et al. 1981; Stevenson 1984; Vargas 1998; Vargas et al. 1998). Examining the record at the scale of the landscape has provided insight into variability in settlement and subsistence patterns (e.g., Vargas et al. 2006). Part of the emphasis on landscapes is a consequence of the practical and conceptual difficulty of working at smaller scales, particularly relevant given the problematic notion of “site” (Dunnell 1992). It is clear to most researchers that the archaeological landscape of Rapa Nui exemplifies the impracticality of the “site” as a unit of observation, recording, and analysis. At least in some domains, the “site” has been regarded as a discrete archaeological unit with ethnographic connotations of “community” (Willey and Phillips 1958), as has been com-

mon in Polynesian archaeology. On Rapa Nui, however, with a nearly continuous distribution of artifacts along scales ranging from small objects to monumental architecture spread over the surface of the island, any identification of “site” from any scale of observation remains decidedly arbitrary.

Unfortunately, even though research has tended to emphasize landscape scale patterning and distributions, the units of observation have focused on intuitive ethnographic or “*emic*” classes (Commendador 2005). These classes (e.g., *umu*, *hare moa*, *hare paenga*), while useful shorthand for communicating general observations, tend to compress variability in typological units, idealize forms, and render meaningful analysis of these entities difficult, particularly outside of an ethnographic context. This problem can be avoided through an *analytic* approach to the functional and stylistic structure of the archaeological record of Rapa Nui. Such an analytic approach examines patterning at multiple scales – artifacts, their aggregations, and multiple aggregations – without the imposition of the “site,” variously conceived. “Non-site” analytic approaches have been successfully demonstrated in other contexts (e.g., Dunnell 1983; Dunnell and Campbell 1977; Ebert 1992), and are certainly applicable to our examinations of the archaeological phenomena of Rapa Nui.

SATELLITE IMAGES

In top-down archaeological research design, satellite images provide an initial means for generating information on the archaeological record over the largest areas. Over the past several decades, satellite imagery has become a powerful and efficient means for documenting the structure of the earth’s surface over large areas. The potential of satellite images in archaeological research, however, is only beginning to be recognized. Part of the slow rate of adoption of satellite-based research in archaeology has been the relatively high cost, the technological knowledge required, and the comparatively low-resolution of images available from early satellites. Thus, the earliest uses of satellite images were limited mostly to remote sensing specialists and the study of landscapes or large archaeological features (e.g., Allan and Richards 1983; Custer 1986; Ebert 1980; Findlow and Confield 1980; Schalk and Lyons 1976).

The past several years, however, have seen an explosion of new sources of satellite images with increasing resolution integrated into user-friendly tools. These new sources include declassified military imagery and the establishment of commercial firms that have launched their own satellites. A new generation of satellites such as Corona, SPIN-2, Orbview-3/4, SPOT, EROS, and Ikonos provide high resolution images¹ that are well below 10 m and available at low cost. Since these images are capable of resolving features of archaeological interest (e.g., structures, deposits), their use has spawned a wide variety of applications in archaeologi-

¹ In general, “high resolution imagery” is defined as a representation of the Earth’s surface of less than 10 m² (Forte 2001:132).

cal research (e.g., Failmezger 2001; Fowler, 1996, 2002; Kennedy 1998, Kouchoukos 2001; Lipo and Hunt 2005; Mumford and Parcak 2002; Philip et al. 2002).

Significantly, the resolution of recent satellite images has proven to be sufficient to resolve linear prehistoric features such as roads (e.g., Ur 2003; Sever and Wagner 1991) and even footpaths (McKee et al. 1994; Sheets and Sever 1991; Sheets 2003). As we showed recently (Lipo and Hunt 2005), it is possible to detect features related to Rapa Nui statue (*moai*) roads provided one can acquire images that resolve features ca. 2 to 5 m across and if views of *moai* road related-features are not obstructed by vegetation or other kinds of ground cover. When these conditions are met, it is possible to construct a map of the prehistoric roads and use satellite images as a primary means to study at least some portion of the paths across which the *moai* were moved (Lipo and Hunt 2005). As the readers of this journal will appreciate, the sparse vegetation of Rapa Nui provides an ideal location for an array of remote sensing applications.

In our initial work using satellite data, we acquired images from DigitalGlobe's QuickBird satellite. This commercial satellite was launched in 2001, orbits the earth every 93.5 minutes and revisits its path every 1 to 3.5 days depending on latitude. Remarkably, the QuickBird satellite is capable of generating panchromatic images with resolutions of 61 to 72 cm and multi-spectral images with resolutions of 244 to 288 cm. Differences in resolutions depend on the degree to which the satellite is off from the nadir when the image is taken. Data in the DigitalGlobe images are geo-processed so that points and features can be located with an accuracy of 23 m at 90% circular error.

The resolution of the QuickBird satellite provides images that are comparable with aerial photographs typically taken at high elevations. Studies have shown that QuickBird imagery is of sufficient resolution to provide a base for mapping between 1 in = 200 ft and 1 in = 400 ft scale (Nale 2002). One of the advantages of QuickBird is its ability to acquire data at near nadir and that the corresponding digital imagery may be considered true orthophotos. An additional advantage that QuickBird has over aerial photographs is that images are available in color, panchroma, and 4-band multi-spectra ranging from blue to near-infrared (400 nm – 900 nm). This multi-spectral information, though at a lower resolution (ca. 2.4 m), is of sufficient quality to provide for a broad range of vegetation and environmental information. Natural color imagery can provide crop, forest, and wetland information. Remarkably, all of these datasets are available for areas across most of the Earth for about US\$30 per square kilometer.

GOOGLE EARTH IMAGERY FOR RAPA NUI

One recently available means of displaying data in a spatially integrative fashion is Google Earth². Google Earth

² <http://earth.google.com/>

³ This imagery is regularly added to and updated by Google.




Figure 2. Satellite image of Rapa Nui (Image 2007 NASA).

mapping service is a free application for browsing and viewing maps, imagery and data from around the world. Utilizing an intuitive graphical interface, images and data are streamed over the Internet to the application as one zooms and “flies” across the planet. The application is impressive and sports imagery across the entire Earth. One particularly useful feature of Google Earth is that for many areas high-resolution color satellite imagery is available³. In November 2006, Google added imagery for Rapa Nui with one-meter resolution (Figure 2). Although a small area of the northern edge of the island is missing and some small portions of the island are covered by clouds, this imagery represents the most complete and highest-resolution coverage for the island that is currently publicly (and *freely*) available. The images provide a remarkable source of information on the environment, landscape, land-use, and the archaeological record, clearly displaying architectural features, *ahu*, paths of *moai* roads and the position of *moai*, *manavai* (agricultural enclosures), and even concentrations of agricultural rock mulch (e.g., see Bork et al. 2004; Wozniak 1999), among many other features. Figures 3 to 9 provide examples of features visible on Rapa Nui in images from Google Earth. We also note that high-resolution images are available for many Pacific Islands (see Figure 10, a view of the Maeva area of Huahine Island, Society Islands, including stone-constructed fish traps).

In our continuing field research addressing multiple aspects of Rapa Nui prehistory we have integrated commercially available satellite images, free images from Google Earth, kite- and blimp-based low elevation aerial photography with pedestrian surveys, mapping, geophysical techniques, and small-scale excavations (e.g., Ayala et al. 2005; Lipo and Hunt 2005; Lipo et al. 2005). This top-down approach is inspired by the conservation ethic that we outline in this paper. Figures 11 and 12 illustrate the correspondence of Google Earth imagery and the higher resolution counterparts of kite/blimp photography of the same areas of



Figure 3. Satellite image of Rano Raraku statue (*moai*) quarry (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 4. Satellite image of Puna Pau Crater red-scoria top-knot (*pukao*) quarry; the *pukao* are visible (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 5. Satellite image of Akahanga, southern coast; the coastal *ahu*, modern roads, stone mulch areas, and other features are visible (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 6. Satellite image of an interior area of Akahanga where multiple *manavai* (agricultural enclosure) clusters are visible (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 7. Satellite image of an interior area of Akahanga where *moai* road features, historic *pirca* (walls), and modern agricultural fields are visible (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 8. Satellite image of Hanga Ho'onu, northeast coast; several large *ahu* structures are visible (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 9. Satellite image of an interior area near Maunga Te Kahu Rere, south coast, where land-use changes in developing *parcelas* for agriculture are visible (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 10. Satellite image of Maeva area of Huahine Island, Society Islands, French Polynesia; the *marae* complex and traditional fish traps are visible (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



Figure 11. Satellite image of an interior area of Oroi, south coast; the *moai* road, *hare paenga*, *manavai*, and other features are visible, but note the greater detail in the kite photograph show in Figure 12 (from Google Earth Mapping Service, <http://earth.google.com>, Image © 2007 DigitalGlobe, Image © 2007 NASA).



12. A kite-based aerial photograph of archaeological features in the interior area of Oroi, south coast; the higher resolution clearly shows the *moai* road, *hare paenga*, *manavai*, and other features, but in greater detail than in the satellite image (Figure 11). Low elevation photography from a kite- or blimp-platform provides details useful in documenting the archaeological record.

the south coast (near Ahu Oroí) of Rapa Nui. For complex, fragile, and sensitive architectural features such as *ahu*, the integration of remote imagery from satellites and kites provides unmatched data in producing detailed maps. Such maps could be produced with little, if any, direct contact with the structure (e.g., walking over it). As preservation concerns increase with continuing impacts, particularly in the remarkable archaeological record of Rapa Nui, these non-invasive mapping techniques will likely prove essential.

We suggest that field data can be mapped and provided in its spatial context using Google Earth and displaying data, such as for *moai*, *ahu*, *moai* roads, and other archaeological features for sharing using Internet venues, as we have developed with the Museo Antropológico P. Sebastián Englert (e.g., Torres Hochstetter et al. in press). Public access and sharing of data – for the Rapanui and research communities in particular – have become particularly urgent with rapid land-use changes on the island.

The recently and freely available high-resolution satellite images from Google Earth should prove useful to researchers working on Rapa Nui. Integration of these sources, no longer just economical, but now *free*, facilitates a top-down research strategy and a conservation ethic. It should also prove to be a vital tool in integrating and sharing archaeological data for purposes of research and preservation (see Kintigh 2006).


REFERENCES

- Allan, J. A., J. S. Latham and R. W. Smith. 1980. Monitoring land and water use in the Gefara region by satellite imagery. *Annual Report for Libyan Studies* 12:79-80.
- Allan, J. A. and T. S. Richards. 1983. Use of satellite imagery in archaeological surveys. *Annual Report for Libyan Studies* 14:4-8.
- Ayala, I., C. Lipo, and T. Hunt. 2005. The use of satellite imagery to study prehistoric agricultural features (*manavai*) on Rapa Nui. *VI International Conference on Rapa Nui and the Pacific*, C. M. Stevenson, J. M. Ramírez, F. J. Morin, and N. Barbacci, eds.: 113-123. Los Osos: Easter Island Foundation.
- Bork, H-R., A. Mieth & B. Tschochner. 2004. Nothing but stones? A review of the extent and technical efforts of prehistoric stone mulching on Rapa Nui. *Rapa Nui Journal* 18:10-14.
- Commendador, A. 2005. *Measuring variability in prehistoric stone construction on Rapa Nui, Chile*. Unpublished M.A. thesis. Department of Anthropology, University of Hawai'i, Manoa.
- Cristino, C., P. Vargas Casanova and R. Izaurieta. 1981. *Atlas Arqueológico de Isla de Pascua*. Centro de Estudios Isla de Pascua, Universidad de Chile. Santiago, Chile: Corporación Toesca.
- Custer, J. F. 1986. Application of LANDSAT data and synoptic remote sensing to predictive models for prehistoric archaeological sites: an example from the Delaware coastal plain. *American Antiquity* 51(3):572-588.
- Dunnell, R. C. 1982. Science, social science, and common sense: The agonizing dilemma of modern archaeology. *Journal of Anthropological Research* 38:1-25.
- Dunnell, R. C. 1983. Aspects of the spatial structure of the Mayo site (15-JO-14), Johnson County, Kentucky. *Anthropological Papers-Museum of Anthropology, University of Michigan* (72):109-165.
- Dunnell, R. C. 1984. The Ethics of Archaeological Significance Decisions. *Ethics and Values in Archaeology*. E. Green, ed.: 61-74. New York: Free Press.
- Dunnell, R. C. 1992. The Notion Site. *Space, Time and Archaeological Landscapes*, J. Rossignol and L. Wandsnider, eds.: 21-41. New York: Plenum Press.
- Dunnell, R. C. & S. K. Campbell. 1977. *Aboriginal Occupation of Hamilton Island, Washington*. Seattle: University of Washington, Dept. of Anthropology, Reports in Archaeology, 4.
- Dunnell, R. and W. Dancey. 1978. Assessment of Significance; cultural resource management plans. *ASCA Bulletin* 5:2-7.
- Ebert, J. I. 1977. The role of remote sensing. *Settlement and Subsistence along the Lower Chaco River*, C.A. Reher, ed.: 191-215. Albuquerque: University of New Mexico Press.
- Ebert, J. I. 1992. *Distributional Archaeology*. Albuquerque: University of New Mexico Press.
- Ebert, J. I. and T. R. Lyons. 1980. Prehistoric irrigation canals identified from Skylab III and Landsat imagery in Phoenix, Arizona. *Cultural Resources Remote Sensing*. T. R. Lyons and Frances J. Mathien, eds.: 209-228. Washington, D.C.: Cultural Resources Management Division, National Park Service.
- Failmezger, V. 2001. High resolution aerial color IR, multi-spectral, hyper-spectral, and SAR imagery over the Oatlands Platanon archaeological site near Leesburg – a pilot study of the new high resolution satellite imagery. *Remote Sensing in Archaeology*. S. Campana and M. Forte, eds.: 143-148. Florence: All'Insegna Del Giglio.
- Findlow, F. J. and L. Confeld. 1980. LANDSAT imagery and the analysis of archaeological catchment territories: a test of the method. *Anthropology* 10 (1):31-52.
- Fowler, M. J. F. 1996. High-resolution satellite imagery in archaeological application: a Russian satellite photograph of the Stonehenge region. *Antiquity* 70 (269):667-671.
- Fowler, M. J. F. 2002. Satellite remote sensing and archaeology: a comparative study of satellite imagery of the environs of Figsbury Ring, Wiltshire. *Archaeological Prospection* 9 (2):55-69.
- Hunt, T. L., C. P. Lipo and S. Sterling. 2001. Posing questions for a scientific archaeology. *Posing Questions for a Scientific Archaeology*. T. L. Hunt, C. P. Lipo and S. Sterling, eds.: 1-21. London: Bergin and Garvey.
- Kennedy, D. 1998. Declassified satellite photographs and archaeology in the Middle East: Case studies from Turkey. *Antiquity* 72(277):553-561.
- Kintigh, K. 2006. The promise and challenge of archaeological data integration. *American Antiquity* 71:567-578.
- Kouchoukos, N. 2001. Satellite images and Near Eastern landscapes. *Near Eastern Archaeology* 64 (1):80-91.
- Lipe, W. D. 1974. A Conservation Model for American Archaeology. *The Kiva* 39:213-248.
- Lipo, C. P. and T. L. Hunt. 2005. Mapping prehistoric statue roads on Easter Island. *Antiquity* 79:158-168.
- Lipo, C. P., T. L. Hunt, and S. Rapu. 2005. Recent geophysical and archaeological studies at 'Anakena. *VI International Conference on Rapa Nui and the Pacific*. C. M. Stevenson, J. M. Ramírez, F. J. Morin, and N. Barbacci, eds.: 97-105. Los Osos: Easter Island Foundation.
- McCoy, P. 1976. *Easter Island Settlement Patterns in the Late Prehistoric and Proto-historic Periods* Bulletin 5. New York: Easter Island Committee, International Fund for Monuments.

- McKee B. R., Sever T. L., and Sheets P. 1994. Prehistoric footpaths in Costa Rica: Remote sensing and field verification. *Archaeology, volcanism and remote sensing in the Arenal region, Costa Rica*. P. Sheets and B. McKee, eds. :142-157. Austin: University of Texas.
- Moratto, M. and M. S. Kelley. 1977. Significance in Archaeology. *The Kiva* 42:193-202.
- Mumford, G. and S. Parcak. 2002. Satellite image analysis and archaeological fieldwork in El-Markha plain (south Sinai). *Antiquity* 76 (294):353-4.
- Nale, D. K. 2002. *QuickBird aerial product comparison*. Prepared by eMap International for Digital Globe, Inc. <http://turimage.net/geospatial/Sci_documents/Aerial_vs_DG.pdf>
- Philip, G., D. Donoghue, A. Beck and N. Galiatsatos. 2002. CORONA satellite photography: an archaeological application from the Middle East. *Antiquity* 76(291):108-118.
- Schalk, R. F. and T. R. Lyons. 1976. Ecological applications of LANDSAT imagery in archeology: an example from the San Juan Basin, New Mexico. *Reports of the Chaco Center* (1):173-186.
- Sever T. L. and Wagner D. W. 1991. Analysis of prehistoric roadways in Chaco Canyon using remotely sensed digital data. *Ancient road networks and settlement hierarchies in the New World*. C. Trombold, ed. :42-52. Cambridge: Cambridge University Press.
- Sheets, P. D. 2003. Prehistoric human footpaths in Costa Rica indicate intimate ties with villages, cemeteries. *ScienceDaily*, October 8, 2003. <<http://www.sciencedaily.com/releases/2003/10/031008064657.htm>>.
- Sheets, P. D. and T. L. Sever. 1991. Prehistoric footpaths in Costa Rica: Transportation and communication in a tropical rainforest. *Ancient Road Networks and Settlement Hierarchies in the New World*, C. D. Trimbald, ed. :53-65. Cambridge: Cambridge University Press.
- Stevenson, C. M. 1984. *Corporate Descent Group Structure in Easter Island Prehistory*. Ph.D. Dissertation, The Pennsylvania State University.
- Torres Hochstetter, F., S. Rapu Haoa, C. Lipo, and T. Hunt. In press. A public database of archaeological resources on Easter Island (Rapa Nui) using Google Earth. *Latin American Antiquity*.
- Ur, J. 2003. CORONA satellite photography and ancient road networks: a northern Mesopotamian case study. *Antiquity* 77 (295):96-115.
- Vargas C., P. 1998. Rapa Nui Settlement Patterns: Types, function, and spatial distribution of households structural components. *Easter Island and East Polynesian Prehistory*, P. Vargas Casanova, ed. :111-130. Santiago: Instituto de Estudios Isla de Pascua, Facultad de Arquitectura y Urbanismo, Universidad de Chile.
- Vargas C., P., C. Cristino and R. Izaurieta. 2006. *1000 años en Rapa Nui: Arqueología del asentamiento*. Santiago: Instituto de Estudios Isla de Pascua, Universidad de Chile.
- Vargas C., P., R. Izaurieta, C. Cristino and C. Arias. 1998. Rapa Nui settlement and subsistence pattern studies on Easter Island. A GIS based analysis of the Archaeological Survey Data. *Easter Island and East Polynesian Prehistory. Proceedings of the Second International Congress on Easter Island and East Polynesia*, P. Vargas, ed. :147-152. Santiago: Universidad de Chile, Instituto de Estudios Isla de Pascua.
- Wiley, G. R. and P. Phillips. 1958. *Method and Theory in American Archaeology*. Chicago: University of Chicago Press.
- Wozniak, J. A. 1999. Prehistoric horticultural practices on Easter Island – lithic mulched gardens and field systems. *Rapa Nui Journal* 13:95-99.



PREHISTORIC RAPA NUI. LANDSCAPE AND SETTLEMENT ARCHAEOLOGY AT HANGA HO'ONU



By CHRISTOPHER STEVENSON & SONIA HAOA

with contributions by Joan Wozniak,
Helene Martinsson-Wallin, & Paul Wallin

As the authors of this book show, contrary to past perceptions, the Easter Island landscape was a highly transformed and managed agricultural terrain that emerged in response to deforestation by the Polynesians who settled there. This volume adds a new dimension to scholarly investigations about why the island's prehistoric society evolved the way it did.

Soft cover, 297 pages & fold-out map • an EIF publication, 2008 [#SH08]
ISBN 978-1-880636-26-8 • PRICE: \$30.00 + shipping